

Dragonflies and damselflies of a montane tropical rainforest in Papua New Guinea

Steffen Oppel

Wildlife Conservation Society, PO Box 277, Goroka, EHP, Papua New Guinea
 steffen.oppel@gmail.com
 http://www.steffen-oppel.com



Introduction

The odonate fauna of Papua New Guinea (PNG) is poorly studied and distribution and habitat requirements of many species are unknown. The human population in PNG is growing rapidly, which potentially results in habitat degradation and a loss of biodiversity.

In this study I surveyed odonates for several months in an aseasonal rainforest in order to compare the odonate communities of a pristine forest and a modified garden landscape. The principal aim of this investigation was to examine which assemblages are at the highest risk from human induced habitat modification.




Study Area

The study area was located in the Crater Mountain Wildlife Management Area (CMVMA), a 2700 km² tract of continuous rainforest in Papua New Guinea (Fig. 1). The climate is equal throughout the year, and both rainfall and temperatures do not exhibit distinctive seasons. I carried out studies at two research stations within the CMVMA, one situated in primary forest, and one in a typical village. The primary study site (Crater Mountain Biological Research Station - CMBRS) is in the centre of the CMVMA, ranging from 850 m to 1300 m a.s.l. Rainfall is abundant (6.5m, Wright et al. 1997), numerous streams and creeks are present, and the area has never been logged. The village of Herowana lies 10 km to the east of CMBRS at an elevation of 1300 m, and consists of mostly subsistence gardens, small coffee plantations and scattered huts. Native forest is still present, but has been degraded by cutting, and there are more open habitats than at CMBRS.


Methods

Dragonflies were sampled between November 2003 and October 2004 on 112 days at CMBRS and on 36 days in Herowana. Specimens were collected with a net and later identified by J. Michalski (Morristown, USA) and N. Donnelly (Binghamton, USA). I used rank-abundance curves, Simpson's diversity index and evenness to describe the community composition at each site. The similarity of both communities was compared with a new Jaccard-type abundance index, calculated in the program EstimateS 7.5 (Colwell 2004, Chao et al. 2005).

At CMBRS I recorded the following habitat variables at every point where an adult odonate was observed: stream order, nature of water body, shading, substrate presence of boulders > 1m, presence of standing pools, presence of floating vegetation, and distance to the nearest sunny patch. The habitat variables recorded were used to classify the odonate assemblage into several groups, representing distinct communities per habitat type. I used a hierarchical cluster analysis to group species into clusters with similar environmental characteristics. I then recorded the use of habitat types in Herowana and at CMBRS, and calculated the percentages of shared and locally endemic species that preferred and occurred in each habitat type.



Procordulia heppdli, the most common dragonfly at a pond in the pristine rainforest at CMBRS.



I trained local assistants from the village of Herowana to assist me in capturing dragonflies.

This resulted in an increase of awareness for dragonflies in the local community.

Results

Community comparison

I recorded a total of 78 species from 13 families in both study areas together, with 21 species occurring both in Herowana and at CMBRS (Table 1). The natural rainforest site had a higher number of species, and a higher diversity and evenness (Fig. 1). The Jaccard abundance index for both communities was 0.477, and differed widely between families.

Ordination of odonate assemblages at CMBRS

The hierarchical cluster analysis identified seven ecologically significant groups (Fig. 2). The position of cluster centroids in relation to environmental variables is presented in Table 2. The group including species associated with rivers contained the highest number of species (20), followed by the assemblage of smaller and shadier temporary creeks with 13 species. Taken together, the assemblages of temporary water sources and the forest interior comprised 39% of all species present at CMBRS. Anisopterans occurred only in four of the assemblages, and only two species occurred in temporary water sources.

Comparison of habitat use and preference

In Herowana, the most species-rich habitat types were ditches, small permanent creeks, and open sunny areas. These three habitats also served as preferred habitat for 68% of all species (Fig. 3). At CMBRS the distribution of preferred habitats was more equal than in Herowana, with five habitat types being equally important (Fig. 3). They served as preferred habitat for 79% of all species. At CMBRS, only one (13%) of those species preferring the forest interior, and three of the nine species preferring small temporary creeks (33%) also occurred in Herowana. In Herowana, more than half of the species preferring open sunny areas, the forest interior, sunny puddles, small permanent creeks and ditches occurred also at CMBRS.

Table 1. Community statistics of the two odonate communities in Crater Mountain Wildlife Management Area, Papua New Guinea, in 2004.

| | Herowana | CMBRS |
|--------------------------------------|----------|----------|
| Number species | 38 | 61 |
| Number families | 10 | 13 |
| Anisoptera | 9 (24%) | 16 (27%) |
| Zygoptera | 29 (76%) | 45 (73%) |
| species total | 78 | |
| species co-occurring | 21 | |
| species only at that site | 17 | 40 |
| Jaccard-Abd Index (Chao et al. 2005) | 0.447 | |
| Simpson | 8.59 | 11.48 |
| Evenness | 0.715 | 0.804 |




Figure 1. Rank-abundance curves for the odonate communities at CMBRS and in Herowana, Crater Mountain Wildlife Management Area, Papua New Guinea, in 2004. Note that for simplicity only 38 species are shown at CMBRS.




Figure 2. Dendrogram of the hierarchical cluster analysis of odonate species at CMBRS, Papua New Guinea, in 2004. The dashed vertical line indicates the 25% cut-off of the maximum squared Euclidean distance to determine clusters. Cluster number is indicated on the left. See Table 2 for environmental characteristics of clusters.

Table 2. Central values of environmental variables for the seven odonate assemblages identified by the cluster analysis in a lower montane tropical rainforest in Papua New Guinea.

| Cluster | number species | class | water speed | nature of water source | shade (%) | substrate | boulder | pools | floating vegetation | mean dist. to sun (m) |
|---------|----------------|----------|-------------|------------------------|-----------|-----------|---------|-------|---------------------|-----------------------|
| 1 | 10 | pond | 0 | permanent | standing | 42 | mud | no | yes | 0.00 |
| 2 | 20 | river | > 1 m/s | permanent | running | 58 | gravel | yes | no | 2.00 |
| 3 | 13 | stream | 0.3 m/s | temporary | running | 88 | mud | no | yes | 7.80 |
| 4 | 5 | creek | 0.7 m/s | temporary | running | 64 | sand | no | yes | 3.60 |
| 5 | 4 | stream | 0.1 m/s | permanent | running | 86 | sand | no | no | 11.40 |
| 6 | 2 | no water | 0 | none | no water | 68 | mud | no | no | 0.00 |
| 7 | 5 | puddle | 0 | temporary | standing | 79 | mud | no | no | 6.20 |

Discussion

The odonate fauna of the Crater Mountain Wildlife Management Area is species rich, and distinctly different from lowland areas in PNG (Mack 1998). This indicates a high level of local endemism on New Guinea, similar to other areas in South East Asia (Orr 2004). Since local endemism renders odonates vulnerable to extinction (Korkeamäki & Suhonen 2002), the conservation of primary habitats in all parts of the island is an essential prerequisite for the preservation of odonate diversity on New Guinea.

The modified forest in Herowana had less species and a more skewed community composition than the natural rainforest at CMBRS. This indicates that the modification of the rainforest results in an unbalanced species assemblage, where a few generalist species benefit from large open areas and dominate the odonate community (Samways & Steyler 1996, Stewart & Samways 1998, Clausnitzer 2003). These generalist species were mostly associated with open and artificial habitats, and were much less common or absent in the natural rainforest.

Species associated with temporary puddles in the forest, small temporary creeks, and the forest interior comprised 39% of the total odonate community at CMBRS. Most of the forest interior and small temporary water species at CMBRS did not occur in Herowana, which suggests that species depending on a closed canopy and temporary water sources are most susceptible to forest modification (Clausnitzer 2003). Therefore, modification of the forest caused by human subsistence gardening may lead to a loss of c. 25% of the species associated with closed canopy rainforest habitats. This highlights the urgent need for the conservation of natural rainforests.

References

Chao, A., Chao, R. J., Colwell, R. K. & Shen, T. L. (2005) A new statistical approach for assessing similarity of species composition with incidence and abundance data. *Ecology Letters*, 8, 883-890.

Clausnitzer, V. (2003) Dragonfly communities in coastal habitats of Kenya: indication of biotope quality and the need of conservation measures. *Biodiversity and Conservation* 12, 333-366.

Colwell, R. K. (2004) EstimateS: statistical estimation of species richness and shared species from samples. *Y.S. University of Connecticut, Storrs, CT, Available at: http://beta.ces.cornell.edu/estimates/*

Dijkstra, A. B. & Stålman, M. A. (2002) Odonate assemblages of running waters in the Upper Guinean forest. *Archiv für Hydrobiologie* 157, 397-412.

Korkeamäki, B. & Suhonen, J. (2002) Diversity and habitat specialization of species affect local extinction in dragonfly Odonata populations. *Ecography* 25, 469-485.

Mack, A. L. (1998) A biological assessment of the Milne Bay Basin, Papua New Guinea (Rural Working Paper 9). Conservation International, Washington, D.C.

Orr, A. G. (2004) Global patterns of odonate in Malaysia, Indonesia, Singapore, and Brunei. *International Journal of Odonatology* 7, 371-384.

Rossi, R. J. (2004) Conservation of Odonata in the South Pacific and Australasia. *International Journal of Odonatology* 7, 139-147.

Stewart, M. J. & Steyler, M. S. (1996) Dragonfly (Odonata) distribution patterns in urban and forest landscapes, and recommendations for riparian management. *Biological Conservation* 78, 279-288.

Stålman, M. A. & Samways, M. J. (1998) Conserving dragonfly (Odonata) assemblages relative to river dynamics in an African savanna game reserve. *Conservation Biology* 12, 661-669.

Wright, D., Janson, H. J., Burke, D. & de Silva-Gunatillake, H. G. (1997) Tree and fauna enumeration and diversity in a one-hectare plot in Papua New Guinea. *Biotropica* 29, 280-280.

Acknowledgements

This study was initiated with the aid of the Wildlife Conservation Society (WCS), and was supported by the International Dragonfly Fund (IDF) and the Worldwide Dragonfly Association (WDA). The Department of Environment and Conservation (DEC) of Papua New Guinea gave the generous permission to collect and export sample specimens of dragonflies and damselflies. I am very grateful to Dr. A. Mack, Dr. D. Wright, and R. Sinclair (WCS) for assistance and logistical support while doing the fieldwork. S. Richards and A. C. Orr shared many valuable thoughts and thus improved the design and analysis of the study. I would also like to thank V. Kalkman, A. Chovanec, V. Sohni, and J. J. Baikuana for providing me with literature. Species identification would have been impossible without the help of J. Michalski and N. Donnelly, and I greatly appreciate their effort.

Fieldwork was carried out with the permission of the Pawan and Gimi people inhabiting the two study areas. I am very grateful to both groups for being supportive and hosting me on their land. My assistants Oliver, Wesley, Joris, and Amoli from the village of Herowana deserve special thanks for their great skill and tireless effort in collecting specimens. I would also like to thank all garden owners in Herowana for allowing me and my assistants to work on their ground.




Figure 3. Proportion of all species preferring a particular habitat type at CMBRS (black columns) and in Herowana (white columns), Crater Mountain Wildlife Management Area, Papua New Guinea, in 2004. Habitat types: (1) open sunny areas without water, (2) forest interior without water, (3) temporary puddles in the forest interior, (4) temporary puddles in open sunny areas, (5) fish-free permanent pond, (6) large open river, (7) small partially shaded river, (8) small, permanent, mostly shaded creek, (9) small temporary creek in the forest, and (10) artificial ditch.



Nososticta finisterra (Protoneuridae, left) and *Rhinophya linza* (Chlorocyphidae, right) were common representatives of both the village and the natural forest communities. Photo courtesy of Stuart Wilson.

This study was supported by the

Wildlife Conservation Society, Papua New Guinea Program
 International Dragonfly Fund
 Worldwide Dragonfly Association

<http://www.wcs.org/international/Asia/175994>
<http://www.calopteryx.de/idf/index.html>
<http://powell.colgate.edu/wad/dragonfly.htm>

